





PubMed	Nucleotide	Protein	Genome	Structu	re Pop	Set Taxono	omy OMIM	
Search PubMe	ed v for	Andrew Andrews Andrews	***	* :	20 14 182 144 14 2 -	Go	Clear	
 	Limits	Preview/Index		istory	Clipboard			
- 	Display	Abstract	Save	Text	Order	Add to Clipbo	ard	
Entrez								
PubMed	□1: J Exp Med 1996 Jun 1;183(6):2581-91					Related Articles, Books		

PubMed Services The role of gp130-mediated signals in osteoclast development: regulation of interleukin 11 production by osteoblasts and distribution of its receptor in bone marrow cultures.

Romas E, Udagawa N, Zhou H, Tamura T, Saito M, Taga T, Hilton DJ, Suda T, Ng KW, Martin TJ.

St. Vincent's Institute of Medical Research, University of Melbourne, Victoria, Australia.

Related Resources Interleukin (IL)-11 is a multifunctional cytokine whose role in osteoclast development has not been fully elucidated. We examined IL-11 production by primary osteoblasts and the effects of rat monoclonal anti-mouse glycoprotein 130 (gp130) antibody on osteoclast formation, using a coculture of mouse osteoblasts and bone marrow cells. IL-1, TNF alpha, PGE2, parathyroid hormone (PTH) and 1 alpha,25-dihydroxyvitamin D3 (1 alpha,25(OH)2D3) similarly induced production of IL-11 by osteoblasts, but IL-6, IL-4, and TGF beta did not. Primary osteoblasts constitutively expressed mRNAs for both IL-11 receptor (IL-11R alpha) and gp130. Osteotropic factors did not modulate IL-11R alpha mRNA at 24 h, but steady-state gp130 mRNA expression in osteoblasts was upregulated by 1 alpha,25(OH)2D3, PTH, or IL-1. In cocultures, the formation of multinucleated osteoclast-like cells (OCLs) in response to IL-11, or IL-6 together with its soluble IL-6 receptor was dose-dependently inhibited by rat monoclonal anti-mouse gp130 antibody. Furthermore, adding anti-gp130 antibody abolished OCL formation induced by IL-1, and partially inhibited OCL formation induced by PGE2, PTH, or 1 alpha,25(OH)2D3. During osteoclast formation in marrow cultures, a sequential relationship existed between the expression of calcitonin receptor mRNA and IL-11R alpha mRNA. Osteoblasts as well as OCLs expressed transcripts for IL-11R alpha, as indicated by RT-PCR analysis and in situ hybridization. These results suggest a central role of gp130-coupled cytokines, especially IL-11, in osteoclast development. Since osteoblasts and mature osteoclasts expressed IL-11R alpha mRNA, both bone-forming and bone-resorbing cells are potential targets of IL-11.

PMID: 8676079 [PubMed - indexed for MEDLINE]







PubMed Nu	ucleotide Protein Genome Structure PopSet Taxonomy OMIM							
Search PubMed	for IL-11R antibodies Go Clear							
	Limits Preview/Index History Clipboard							
<u></u>	Display Summary							
ntrez ubMed	Show: 20 T Items 1-10 of 10 One page							
	1: Campbell CL, Guardiani R, Ollari C, Nelson BE, Quesenberry PJ, Savarese TM. Related Articles							
ubMed ervìces	Interleukin-11 receptor expression in primary ovarian carcinomas. Gynecol Oncol. 2001 Feb;80(2):121-7. PMID: 11161848 [PubMed - indexed for MEDLINE]							
	☐2: Prybylowski KL, Wolfe BB. Developmental differences in alternative splicing of the NR1 protein in rat cortex and cerebellum. Brain Res Dev Brain Res. 2000 Oct 28;123(2):143-50. PMID: 11042343 [PubMed - indexed for MEDLINE]							
elated esources	3: Barton VA, Hall MA, Hudson KR, Heath JK. Interleukin-11 signals through the formation of a hexameric receptor complex. J Biol Chem. 2000 Nov 17;275(46):36197-203. PMID: 10948192 [PubMed - indexed for MEDLINE]							
	4: Blanc C, Vusio P, Schleinkofer K, Boisteau O, Pflanz S, Minvielle S, Grotzinger J, Muller-Newen G, Heinrich PC, Jacques Y, Montero-Julian FA. Monoclonal antibodies against the human interleukin-11 receptor alpha-chain (IL-11Ralpha) and their use in studies of human mononuclear cells. J Immunol Methods. 2000 Jul 31;241(1-2):43-59. PMID: 10915848 [PubMed - indexed for MEDLINE]							
	The leukin-11 enhancement of VLA-5 mediated adhesion of CD34+ cells from cord blood to fibronectin is associated with the PI-3 kinase pathway. In Vivo. 2000 Mar-Apr;14(2):331-7. PMID: 10836206 [PubMed - indexed for MEDLINE]							
	6: Jiang H, Sugimoto K, Sawada H, Takashita E, Tohma M, Gonda H, Mori KJ. Related Article Mutual education between hematopoietic cells and bone marrow stromal cells through direct cell-to-cell contact: factors that determine the growth of bone marrow stroma-dependent leukemic (HB-1) cells. Blood. 1998 Aug 1;92(3):834-41. PMID: 9680351 [PubMed - indexed for MEDLINE]							
	7: Chevalier S, Fourcin M, Robledo O, Wijdenes J, Pouplard-Barthelaix A, Gascan H. Interleukin-6 family of cytokines induced activation of different functional sites							

expressed by gp130 transducing protein.
J Biol Chem. 1996 Jun 21;271(25):14764-72.
PMID: 8662918 [PubMed - indexed for MEDLINE]

8: Romas E, Udagawa N, Zhou H, Tamura T, Saito M, Taga T, Hilton DJ, Suda Related Articles T, Ng KW, Martin TJ.

The role of gp130-mediated signals in osteoclast development: regulation of interleukin 11 production by osteoblasts and distribution of its receptor in bone marrow cultures.

J Exp Med. 1996 Jun 1;183(6):2581-91.

PMID: 8676079 [PubMed - indexed for MEDLINE]

9: Tani Y, Nishimoto N, Ogata A, Shima Y, Yoshizaki K, Kishimoto T.

Related Articles

Gp130 in human myeloma/plasmacytoma.

Curr Top Microbiol Immunol. 1995;194:229-33. No abstract available.

PMID: 7895494 [PubMed - indexed for MEDLINE]

10: Fourcin M, Chevalier S, Lebrun JJ, Kelly P, Pouplard A, Wijdenes J, Gascan Related Articles H.

Involvement of gp130/interleukin-6 receptor transducing component in interleukin-11 receptor.

Eur J Immunol. 1994 Jan;24(1):277-80.

PMID: 8020567 [PubMed - indexed for MEDLINE]

Display Summary Save Text Order Details Add to C	lipboard
Show: 20 Items 1-10 of 10	One page.

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer